F 588

88194
F.588

A

SET

OF

ANATOMICAL TABLES,

WITH

EXPLANATIONS,

AND AN

ABRIDGMENT,

OF THE

PRACTICE of MIDWIFERY,

With a View to illuſtrate a

TREATISE

On that SUBJECT,

AND

COLLECTION of CASES.

The SECOND EDITION, Corrected.

By *WILLIAM SMELLIE*, M. D.

LONDON:

Printed in the Year M.DCC.LXI.

PREFACE.

AS in a long courſe of teaching and practice in Midwifery, I hope I may without vanity ſay, that I have done ſomething towards reducing that Art, into a more ſimple and mechanical method than has hitherto been done, I have attempted to explain the ſame in my Treatiſe of the Theory and Practice of Midwifery and Collection of Caſes, and finding that moſt of the repreſentations hitherto given of the parts ſubſervient to Uterine geſtation *and* parturition *were in many reſpects deficient, I have been induced to undertake the following* Tables, *with a view to ſupply in ſome meaſure the defects of others, and at the ſame time to illuſtrate what I have taught and written on the Subject. How far I have obtained thoſe ends, it belongs to others to judge ; I ſhall only beg leave to obſerve here by way of Preface, that the greateſt part of the figures were taken from Subjects prepared on purpoſe, to ſhew every thing that might conduce to the improvement of the young Practitioner, avoiding however the extreme Minutiæ, and what elſe ſeemed foreign to the preſent deſign ; the ſituation of parts, and their reſpective dimenſions being more particularly attended to, than a minute anatomical inveſtigation of their ſtructure.*

As theſe Tables *may poſſibly fall into the hands of ſome who have not ſeen my former work, I have added an abridgment of the Practice, which though far from being complete, may ſerve to illuſtrate ſeveral things which otherwiſe by a bare repreſentation would be hardly intelligible.*

References are made to Vol. I. II. *and* III. *by* Vol. I. *I mean that which I firſt publiſhed in the year* 1752. *and contains a view of the Theory and Practice of Midwifery ;* Vol. II. *is the Collection of Caſes mentioned above juſt publiſhed ; and* Vol. III. *is a continuation of the ſame, and is already far advanced. My firſt plan for theſe* Tables *confined them to the number of Twenty-two, which Mr.* Rymſdyke *had finiſhed above two Years ago ; but I ſoon ſaw that a further illuſtration and conſequently an addition to that number was neceſſary. In eleven of theſe Dr.* Camper, *formerly* Profeſſor of Medicine *at* Franequer *in* Frieſland, *now* Profeſſor of Anatomy *and* Botany *at* Amſterdam, *greatly aſſiſted me,* viz. Table XII. XVI. XVII. XVIII. XIX. XXIV. XXVI. XXVII. XXVIII. XXXIV. *and* XXXVI. *The reſt were drawn by Mr.* Rymſdyke, *except the thirty ſeventh and thirty ninth, which were done by another hand. The whole of the Drawings are faithfully engraved by Mr.* Grignion, *delicacy and elegance however has not been ſo much conſulted as to have them done in a ſtrong and diſtinct manner, with this view chiefly, that from the cheapneſs of the work it may be rendered of more general uſe.*


A
D
D
I
I
I
I
B
C
H
H
I
I
G
G
F
E
E

THE

FIRST TABLE

Represents in a Front-View the Bones of a well formed *Pelvis.*

A the five *Vertebræ* of the Loins.
B The *Os Sacrum.*
C The *Os Coccygis.*
D.D The *Ossa Iliûm.*
E.E The *Ossa Ischiûm.*
F The *Ossa Pubis.*
G The *Foramina Magna.*
H.H The *Acetabula.*
I.I.I.I.I.I The Brim of the *Pelvis*, or that circumference of its cavity, which is described at the sides, by the inferior parts of the *Ossa Iliûm*, and at the back and fore-parts, by the superior parts of the *Ossa Pubis* and *Sacrum.*

In this Table besides the general structure and figure of the several Bones, the dimensions of the Brim of the *Pelvis*, and the distance between the under parts of the *Ossa Ischiûm*, are particularly to be attended to, from which it will appear that the cavity at the Brim is commonly wider from side to side, than from the Back to the fore-part, but that the sides below are in the contrary proportion. The Reader, however, ought not from this to conclude, that every *Pelvis* is similar in figure and dimensions, since even well formed ones differ in some degree from each other. In general, the Brim of the *Pelvis* measures about five inches and a quarter from side to side, and four inches and a quarter from the back to the fore-part; there being likewise the same distance between the inferior parts of the *Ossa Ischiûm.* All these measures, however, must be understood as taken from the Skeleton, for in the Subject, the cavity of the *Pelvis* is considerably diminished by its teguments and contents. Correspondent also to this diminution the usual dimensions of the head of the full grown *Foetus* are but three inches and a half from ear to ear, and four inches and a quarter from the fore to the hind-head.

Vide Tab. XVI. XVII. XVIII. Also Vol. I. Chap. I. Sect. 1. 2. 3. where the form and dimensions of the *Pelvis*, as well as of the head of the *Foetus*, and the manner in which the same is protruded in Labour through the Bason, are fully treated of. Consult likewise Vol. II. Col. 1. N° 1. 2. where Cases are given of complaints of the *Pelvis* arising from difficult Labours.

THE

SECOND TABLE

Gives a lateral and internal View of the *Pelvis*, the ſame being divided longitudinally.

A The three lower *Vertebræ* of the Loins.
B The *Os Sacrum.*
C The *Os Coccygis.*
D The left *Os Iliûm.*
E The left *Os Iſchiûm.*
F The *Os Pubis* of the ſame ſide.
G The acute proceſs of the *Os Iſchiûm.*
H The *Foramen Magnum.*
I.I.I The Brim of the *Pelvis.*

This Plate ſhews the diſtance from the ſuperior part of the *Os Sacrum* to the *Oſſa Pubis*, as well as from the laſt mentioned Bones to the *Coccyx*, which in each amounts to about four inches and a quarter. The depth likewiſe is ſhewn of the poſterior, lateral and anterior parts of the *Pelvis*, not in the line of the Body, but in that of the *Pelvis* from its Brim downward, which is generally three times deeper on the poſterior than anterior part, and twice the depth of the laſt at the ſides.

From this view appears alſo the Angle which is formed by the laſt *Vertebra* of the loins and the ſuperior part of the *Os Sacrum*, as likewiſe the concavity or hollow ſpace in the poſterior internal part of the *Pelvis*, ariſing from the curvature of the laſt mentioned Bone and *Coccyx*; finally, the diſtance from which to the poſterior parts of the *Oſſa Iſchiûm* is here expreſt.

Vide Tab. XVI. XVII. XVIII. XIX. Alſo Vol. I. and II. as referred to in the former Table.

A
D
I
I
I
B
F
G
H
E
C

A
D
D
B
C
H
G
G
H
F
E
E

THE

THIRD TABLE

Fxhibits a Front-View of a diſtorted *Pelvis.*

A the five *Vertebræ* of the Loins.
B The *Os Sacrum.*
C The *Os Coccygis.*
D.D The *Oſſa Iliûm.*
E.E The *Oſſa Iſchiûm.*
F The *Oſſa Pubis.*
G.G The *Foramina Magna.*
H.H The *Acetabula.*

From this Plate may appear the great danger incident to both Mother and Child when the *Pelvis* is diſtorted in this manner; it being only two inches and an half at the Brim from the poſterior to the anterior part, and the ſame diſtance between the inferior parts of each *Os Iſchiûm.* *Vide* Tab. XXVII. where the *Pelvis* is one quarter of an inch narrower at the Brim than this, but ſufficiently wide below. Various are the forms of diſtorted Baſons, but the laſt mentioned is the moſt common. It is a great happineſs however in practice that they are ſeldom ſo narrow, though there are inſtances where they have been much more ſo. The danger in all ſuch caſes muſt encreaſe or diminiſh, according to the degree of diſtortion of the *Pelvis*, and ſize of the Child's head.

Vide Vol. I. Lib. 1. Chap. I. Sect. 4, 5. and Vol. II. Coll. 1. N°. 3, 4, 5. Alſo Coll. 21. 27, and 29.

THE

FOURTH TABLE

Shews the external Female parts of Generation.

A The lower part of the *Abdomen.*
B.B The *Labia pudendi* ſeparated.
C The *Clitoris* and *Præputium.*
D.D The *Nymphæ.*
E The *Foſſa magna,* or *Os Externum.*
F The *Meatus urinarius.*
G The *Frænum Labiorum.*
H The *Perinæum.*
I The *Anus.*
K The part that covers the extremity of the *Coccyx.*
L.L The parts that cover the Tuberoſities of the *Oſſa Iſchiûm.*

As it is of great conſequence to every Practitioner in Midwifery, to know exactly the Situation of the parts concerned in Parturition, and which have not been accurately deſcribed by former Anatomiſts, with a view to this particular Branch; I have given this Draught from one of the preſerved Subjects which I keep by me, in order to demonſtrate theſe parts in the ordinary Courſe of my Lectures. From a View then of the ſituation of the parts it appears, that the *Os Externum* is not placed in the middle of the inferior part of the *Pelvis,* but at the anterior and inferior part of the *Pubes,* and that the *Labia* cover likewiſe the anterior part of theſe bones.

Seoondly, It may be obſerved, that as the *Frænum Labiorum,* which is nearly adjoining to the inferior part of the *Oſſa Pubis,* is only about an inch from the *Anus,* between which and the *Coccyx* there is about three inches diſtance, it follows, that the *Anus* is nearer to the firſt mentioned Bones than to the latter.

Thirdly, The View of this and the following Table will furniſh proper hints, with reſpect to the method of touching or examining the *Os Uteri* without hurting or inflaming the parts, as it appears that the *Os Externum* is placed forwards towards the *Pubes* and the *Os Uteri* backwards towards the *Rectum* and *Coccyx.* By this wiſe Mechaniſm of nature many inconveniencies are often prevented, which muſt happen, if theſe parts were oppoſite to each other, and ſituated in the middle of the inferior part of the *Pelvis,* particularly a *Prolapſus* of the *Vagina* and *Uterus,* either in the unimpregnated ſtate, or in any of the firſt four months of Pregnancy; as alſo too ſudden deliveries in any of the laſt months.

Fourthly, From a view of the Situation of the parts, it will appear, that in Labour, when the *Os Uteri* is ſufficiently opened to allow a paſſage for the Head of the *Foetus,* the ſame is protruded to the lower part of the *Vagina,* by which the external parts are puſhed out in form of a large Tumor, as in Table XV.

Laſtly, It may be obſerved, that when it is neceſſary to dilate the *Os Externum,* the principal force ought to be applied downwards, and towards the *Rectum,* to prevent the *Urethra* and neck of the Bladder from being hurt or inflamed.

Vide Vol. I. Lib. I. Chap. 2. Sect. 1. Vol. II. Coll. 2.


A
B
C
D
E
F
G
H
I
K
L

Fig. 1.



Fig. 2.

Fig. 3.



THE

FIFTH TABLE.

FIGURE I. Gives a front View of the *Uterus* in *Situ* ſuſpended in the *Vagina*; the anterior parts of the *Oſſa Iſchiûm* with the *Oſſa Pubis*, *Pudenda*, *Perinæum*, and *Anus*, being removed in order to ſhew the internal Parts.

A The laſt *Vertebra* of the *Loins*.
B.B The *Oſſa Iliûm*.
C.C The *Acetabula*.
D.D The inferior and poſterior parts of the *Oſſa Iſchiûm*. *Vide* Table XXIX. where the *Oſſa Pubis* and the anterior parts of the *Oſſa Iſchiûm* are repreſented by dotted Lines.
E The Part covering the Extremity of the *Coccyx*.
F The inferior part of the *Rectum*.
G.G The *Vagina* cut open longitudinally and ſtretched on each ſide of the *Collum Uteri*, to ſhew in what manner the *Uterus* is ſuſpended in the ſame.
H.H Part of the *Veſica Urinaria* ſtretched on each ſide of the *Vagina* and inferior part of the *Fundus Uteri*.
I The *Collum Uteri*.
K The *Fundus Uteri*.
L.L The *Tubæ Fallopianæ* and *Fimbriæ*.
M.M The *Ovaria*.
N.N The *Ligamenta lata* and *rotunda*.
O.O The ſuperior part of the *Rectum*.

FIGURE II. Gives a View of the internal Parts as ſeen from the right *Groin*, the *Pelvis* being divided longitudinally.

A The loweſt *Vertebra* of the *Loins*.
B.C The *Os Sacrum* and *Coccyx*, with the Integuments.
D The left *Os Iliûm*.
E The inferior part of the left *Os Iſchiûm*.
F The *Os Pubis* of the ſame ſide.
G The *Foramen Magnum*.
H The *Acetabulum*.
I.I.I The inferior Part of the *Rectum* and *Anus*.
K The *Os Externum* and *Vagina*; the *Os Uteri* lying looſely in the ſame.
L The *Veſica Urinaria*.
M.N The *Collum* and *Fundus Uteri*, with a View of the cavity of both. The attachment of the *Vagina* round the outſide of the Lips of the Mouth of the *Womb* is here likewiſe ſhewn, as alſo the ſituation of the *Uterus*, as it is preſſed downwards and backwards by the *Inteſtines* and *Urinary Bladder*, into the concave, and inferior part of the *Os Sacrum*.
O The *Ligamenta lata* and *rotunda* of the left ſide.
P.P The *Fallopian* Tube with the *Fimbriæ*; and
Q The *Ovarium* of the ſame ſide.
R.R The ſuperior part of the *Rectum*, and inferior part of the *Colon*.

FIGURE III. Gives a front View of the *Uterus* in the beginning of the firſt Month of Pregnancy; the anterior part being removed that the *Embryo* might appear through the *Amnios*, the *Chorion* being diſſected off

A The *Fundus Uteri*.
B The *Collum Uteri*, with a View of the rugous Canal that leads to the Cavity of the *Fundus*.
C The *Os Uteri*.
Vide Vol. I. Lib. I. Chap. 2. Sect. 2, 3. Vol. II. Coll. 3.

THE

SIXTH TABLE.

FIGURE I. In the ſame View and Section of the Parts as in the firſt figure of the former Table, ſhews the *Uterus* as it appears in the ſecond or third Month of Pregnancy, it's anterior part being here likewiſe removed.

F The *Anus*.
G The *Vagina* with it's *Plicæ*.
H.H The poſterior and inferior part of the *Urinary Bladder* extended on each ſide, the anterior and ſuperior part being removed.
I.I The Mouth and Neck of the *Womb*, as raiſed up when examining the ſame by the touch, with one of the fingers in the *Vagina*.
K.K The *Uterus* as ſtretched in the ſecond or third Month, containing the *Embryo*, with the *Placenta* adhering to the *Fundus*.

It appears from this and the former Table that at this time nothing can be known, with reſpect to Pregnancy, from the touch in the *Vagina*, as the reſiſtance of the *Uterus* is ſo inconſiderable that it cannot prevent it's being raiſed up before the finger; and even were it kept down, the length of the Neck would prevent the ſtretching being perceptible. The *Uterus* likewiſe not being ſtretched above the *Pelvis*, little change is made as to the figure of the *Abdomen*, further than that the *Inteſtines* are raiſed a little higher; whence poſſibly the old Obſervation of the *Abdomen* being a little flatter at this Period than uſual from the *Inteſtines* being preſſed more to each ſide. Women at this Period miſcarry oftner than at any other: it is a great happineſs however in Practice that although they are frequently much weakened by large diſcharges, yet they rarely ſink under the ſame, but are ſooner or later relieved by Labour coming on, which gradually ſtretches the Neck and Mouth of the *Womb*, by the *Membranes* being forced down with the Waters, and if the *Placenta* is ſeparated from the internal ſurface of the *Uterus*, all it's contents are diſcharged. But if the *Placenta* ſtill adheres, the *Membranes* break, the Waters and *Eœtus* are expelled, and the flooding diminiſhes from the *Uterus* contracting cloſe to the *Secundines*, which alſo are uſually diſcharged ſooner or later.

From the ſtructure finally of the parts as repreſented in this and the former Table, it may appear that it is much ſafer to reſtrain the Flooding, and ſupport the Patient, waiting with Patience the efforts of Nature, than to endeavour to ſtretch the *Os Uteri* and deliver either with the Hand or Inſtruments, which might endanger a laceration and inflammation of the parts.

Vide C in Table XXXVII. alſo Vol. I. Lib. II. Chap. 2. Sect. 2, 3, 4. Vol. II. Coll. 12. No. 2.

FIGURE II. Repreſents the *Uterus* in the fourth or fifth Month of Pregnancy, in the ſame View and Section of the Parts with the former figure, exepting that in this the anterior part of the *Collum Uteri* is not removed.

In the natural ſituation the Mouth and Lips of the *Womb* are covered with the *Vagina*, and theſe parts are contiguous to each other, but here the *Vagina* G is a little ſtretched from the Neck and Lips of the former in order to ſhew the parts more diſtinctly. I The Neck of the *Womb*, which appears in this figure thicker ſhorter and ſofter than in the former. K The inferior part of the *Fundus Uteri*, the ſtretching of which can ſometimes be felt through the *Vagina* by puſhing up a finger on the anterior or lateral part of the ſame.

The *Uterus* now is ſo largely ſtretched as to fill all the upper part of the *Pelvis* and begins alſo to increaſe ſo much as to reſt on the Brim, and to be ſupported by the ſame, the *Fundus* at the ſame time being raiſed conſiderably above the *Pubes*. From the *Abdomen* being now more ſtretched the Woman is more ſenſible of her growing bigger, and the *Uterus* alſo from the counter-preſſure of the Contents and Parietes of the *Abdomen* is kept down and the *Os Uteri* prevented from riſing before the finger as formerly. In lean Women the ſtretching of the *Uterus* can ſometimes be perceived in the *Vagina* at this period as well as above the *Pubes:* but nothing certain can be diſcovered from the reſiſtance or feel of the Mouth of the *Womb* or *Lips* which are commonly the ſame in the firſt Months of Pregnancy as before it.

The ſize or bulk of the *Fœtus* is finally here to be obſerved with the *Placenta* adhering to the poſterior part of the *Uterus*.

Vide the references to Vol. I. and II. in the former Table.

Fig. 1st



Fig. 2d



C
C
A
C
B
C
A
A
D
D

THE

SEVENTH TABLE

Repreſents the *Abdomen* of a Woman opened, in the ſixth or ſeventh Month of Pregnancy.

A.A.A.A The Parietes of the *Abdomen* opened, and turned back to ſhew.

B The *Uterus*.

C.C.C The *Inteſtines* raiſed upwards.

D The *Labia Pudendi*, which are ſometimes affected in Pregnancy with *Oedematous* ſwellings, occaſioned by the preſſure of the *Uterus* upon the returning Veins, and *Lymphatics*. If the *Labia* are ſo tumefied as to obſtruct the Patient's walking, the complaint is removed by puncturing the parts affected. By which means the ſerous fluid is diſcharged for the preſent; but commonly recurs and the ſame Operation muſt be repeated ſeveral times perhaps before delivery, after which however, the tumefaction entirely ſubſides. Here it may be obſerved that this complaint can ſeldom or never obſtruct delivery, as the *Labia* are ſituated at the anterior part of the *Oſſa Pubis*, and can rarely affect the ſtretching of the *Frænum*, *Perinæum*, *Vagina* and *Rectum*. From this figure it appears that the ſtretching of the *Uterus* can eaſily be felt at this period in lean Subjects, through the Parietes of the *Abdomen*; eſpecially if the *Inteſtines* do not lie before it. In general indeed as the *Uterus* ſtretches, it riſes higher, by which means the *Inteſtines* are likewiſe raiſed higher, and are alſo preſſed to each ſide. Hence the nearer the Woman is to her full time, the ſtretching is the more eaſily felt.

Vide Vol. I. Lib. I. Chap. 3. Sect. 3. Lib. III. Chap. 1. Sect. 2. and Vol. II. Coll. 12, 13.

THE

EIGHTH TABLE

In the ſame View and Section of the parts as in Table VI. is repreſented the *Uterus* of the former Table, in order to ſhew its contents, and the internal parts as they appear in the ſixth or ſeventh Month of Pregnancy.

A The *Uterus* ſtretched up to the *Umbilical* Region.
B.B The ſuperior part of the *Oſſa Ilium.*
C.C The *Acetabula.*
D.D The remaining poſterior parts of the *Oſſa Iſchium.*
E The *Anus.*
F The *Vagina.*
G The Bladder of *Urine.*
H The neck of the *Womb* ſhorter than in Table VI. and raiſed higher by the ſtretching of the *Uterus* above the Brim of the *Pelvis.*
I The Veſſels of the *Uterus* larger than in the unimpregnated ſtate.
K K The *Placenta* adhering to the inferior and poſterior part of the *Uterus.*
L.L The *Membranes* that ſurround the *Fœtus*, the Head of which is here repreſented (as well as of thoſe in Table VI.) ſituated downwards at the inferior part of the *Uterus*, and which I am apt to believe is the uſual ſituation of the *Fœtus*, when at reſt, and ſurrounded with a great quantity of Waters, as the Head is heavier than any other part. With reſpect to the ſituation of the Body of the *Fœtus*, tho' the fore-parts are often turned towards the ſides and poſterior parts of the *Uterus*, they are here as well as in the foregoing Table repreſented at the anterior part or forwards, in order to ſhew them in a more diſtinct, and pictureſque manner.

Vide Vol. I. Lib. I. Chap. 3. Sect. 3, 4. Vol. II. Coll. 13. N°. 1.

From this Table may appear the difficulty of ſtretching the *Os Uteri* in flooding Caſes, even at this period, from the length and thickneſs of the neck of the *Womb*, eſpecially in a firſt Pregnancy: much the ſame method however is to be followed here as was directed in Table VI. 'till Labour comes on to dilate the *Os Uteri.* If the Flooding is then conſiderable, the *Membranes* ſhould be broken, that the *Uterus* may contract, and thereby leſſen the diſcharge. The Labour likewiſe if it is neceſſary, may be aſſiſted by dilating the *Os Uteri* in time of the pains, which alſo, if wanting, may be provoked by the ſame method when the Patient is in danger. If this danger is imminent and the Woman ſeems ready to expire, the *Uterus* as appears from this Table is at this time ſufficiently ſtretched to receive the Operator's hand to extract the *Fœtus*, if the *Os Internum* can be ſafely dilated.

Laſtly, It may be obſerved that Women are in greater danger at this period and afterwards than in the former Months.

Vide Vol. I. Lib. III. Chap. 4. Sect. 3. No. 1, 2, 3. Vol. III. Coll. 33. No. 2. See alſo in the *Edinburgh* Phyſical and Literary Obſervations, Art. XVII. The Diſſection of a Woman with Child, by Dr. *Donald Monro*, Phyſician at *London.*


A
L
B
B
K
K
C
C
I
H
D
E
G
G
F
E

A
L
K
L
B
B
M
I
C
C
D
D
H
G
G
F
E

THE

NINTH TABLE.

In the ſame View and Section of the parts with the former, repreſents the *Uterus* in the eighth or ninth Month of Pregnancy.

A The *Uterus* as ſtretched to near its full extent, with the waters, and containing the *Fœtus* intangled in the *Funis*, the Head preſenting at the upper part of the *Pelvis*.
B.B The ſuperior part of the *Oſſa Iliûm*.
C.C The *Acetabula*.
D.D The remaining poſterior parts of the *Oſſa Iſchiûm*.
E The *Coccyx*.
F The inferior part of the *Rectum*.
G.G.G The *Vagina* ſtretched on each ſide.
H The *Os Uteri*, the lips of which appear larger and ſofter than in the foregoing Table, the neck of the *Womb* being likewiſe ſtretched to it's full extent, or entirely obliterated.
I.I Part of the *Veſica Urinaria*.
K.K The *Placenta* at the ſuperior and poſterior part of the *Uterus*.
L.L The *Membranes*.
M The *Funis Umbilicalis*.

This and the foregoing Table ſhew in what manner the *Uterus* ſtretches, and how it's neck grows ſhorter, in the different periods of Pregnancy; as alſo the magnitude of the *Fœtus* in order more fully to explain Vol. I. Lib. 1. Chap. 3. Sect. 4, 5. alſo Lib. 3. Chap. 1. Sect. 1. 2. likewiſe Vol. II. Coll. 13. No. 1.

Notwithſtanding it has been handed down as an invariable Truth, from the earlieſt accounts of the Art, to the preſent times, that when the Head of the *Fœtus* preſented, the Face was turned to the poſterior part of the *Pelvis*, yet from Mr. *Ould*'s Obſervation as well as from ſome late Diſſections of the *gravid Uterus*, and what I myſelf have obſerved in Practice, I am led to believe that the Head preſents for the moſt part as is here delineated, with one Ear to the *Pubes*, and the other to the *Os Sacrum*; tho' ſometimes this may vary according to the form of the Head, as well as that of the *Pelvis*.

Conſult Dr. *Hunter*'s elegant Plates of the *gravid Uterus*.

THE

TENTH TABLE

Gives a front View of Twins in *Utero*, in the beginning of Labour, the anterior parts being removed, as in the preceding Tables.

A The *Uterus* as ſtretched with the *Membranes* and Waters.
B.B The ſuperior parts of the *Oſſa Iliûm.*
C.C The *Acetabula.*
D.D The *Oſſa Iſchiûm.*
E The *Coccyx.*
F The lower part of the *Rectum.*
G.G The *Vagina.*
H The *Os Internum* ſtretched open about a Finger's breadth with the *Membranes* and Waters in time of Labour-pains.
I.I The inferior part of the *Uterus* ſtretched with the Waters which are below the Head of the Child that preſents.
K.K The two *Placentas* adhering to the poſterior part of the *Uterus*, the two *Fœtus*'s lying before them, one with its Head in a proper poſition, at the inferior part of the *Uterus*, and the other ſituated præternaturally, with the Head to the *Fundus:* the Bodies of each are here entangled in their proper *Funis*, which frequently happens in the natural as well as præternatural poſitions.
L.L.L The *Membranes* belonging to each *Placenta.*

This repreſentation of Twins, according to the Order obſerved in my Treatiſe of Midwifery, ought to have been placed among the laſt Tables; but as that was of no conſequence, I have placed it here in order to ſhew the *Os Uteri* grown much thinner than in the former Figure, a little open, and ſtretched by the Waters and *Membranes* which are puſhed down before the Head of one of the *Fœtus*'s in time of a Labour-pain. With reſpect to the poſition of Twins it is often different in different caſes; but was thus, in a late Diſſection of a *gravid Uterus* by Dr. *Mackenzie.*

Vide Vol. I. Lib. 3. Chap. 1. Sect. 4. and Chap. 5. Sect. 1. and Vol. II. Coll. 14. and Vol. III. Coll. 37.

A
L
L
K
K
B
B
C
C
F
G
D
D
H
G
F
E

A
M
K
B
L
B
C
D
G
H
I
G
F
E
C

THE

ELEVENTH TABLE

Exhibits another front View of the *gravid Uterus* in the beginning of Labour; the anterior parts being removed, as in the former Table; but in this the *Membranes*, not being broken, form a large bag containing the Waters and *Fœtus*.

A The ſubſtance of the *Uterus*.

B.B.C.C.D.D The Bones of the *Pelvis*.

E The *Coccyx*.

F The inferior part of the *Rectum*.

G.G.G.G The *Vagina*.

H.H The Mouth of the *Womb* largely ſtretched in time of a pain, with I the *Membranes* and Waters. This Circumſtance makes it uſually certain that Labour is begun, whereas from the degree of dilatation repreſented in the former Table there is little to be aſcertained, unleſs the pains are regular and ſtrong, the *Os Uteri* being often found more open ſeveral days, and even weeks, before Labour commences.

K The *Chorion*.

L The ſame diſſected off at the inferior part of the *Uterus*, in order to ſhew the Head of the *Fœtus* through the *Amnios*. *N.B.* This Hint is taken from one of Dr. *Albinus*'s Tables of the *gravid Uterus*.

M The *Placenta*, the external convex Surface of which divided into a number of *Lobes*, is here repreſented, it's concave internal part being covered by the *Chorion*.

The *Placenta* has been found adhering to all the different parts of the internal ſurface of the *Uterus*, and ſometimes even over the inſide of the *Os Uteri*; this laſt manner of adheſion however always occaſions Floodings as ſoon as the ſame begins to dilate.

Tables VI. VIII. IX. X. ſhew the internal ſurface of the *Placenta* towards the *Fœtus*, with the Veſſels compoſing it's ſubſtance proceeding from the *Funis* which is inſerted, in different *Placentas* into all the different parts of the ſame, as well as in the middle.

The thirtieth and thirty third Tables ſhew the inſertion of the *Funis* into the *Abdomen* of the *Fœtus*.

With reſpect to the expulſion of the *Placenta*, when the *Membranes* break, the *Uterus* contracts as the Waters are evacuated 'till it comes in contact with the Body of the *Fœtus*: the ſame being delivered, the *Uterus* grows much thicker, and contracts cloſely to the *Placenta* and *Membranes*, by which means they are gradually ſeparated, and forced into the *Vagina*. This ſhews that we ought to follow the method which nature teaches, waiting with patience, and allowing it to ſeparate in a ſlow manner, which is much ſafer practice, eſpecially when the Patient is weak; as the diſcharge is neither ſo great or ſudden as when the *Placenta* is hurried down in the too common method. But then we muſt not run into the other extreme, but aſſiſt, when nature is not ſufficient to expel the ſame.

Vide Vol. I. Lib. III. Chap. 1. Sect. 4. Chap. 2. Sect. 2. 5. Vol. II. Coll. 14. 23.

THE

TWELFTH TABLE

Shews (in a lateral View and longitudinal Diviſion of the Parts) the *gravid Uterus*, when Labour is ſomewhat advanced.

A The loweſt *Vertebra* of the Back.
B The *Scrobiculus Cordis*, the diſtance from which to the laſt mentioned *Vertebra* is here ſhewn by dotted lines; as alſo part of the Region below the *Diaphragm*.
C.C The uſual thickneſs and figure of the *Uterus* when extended with the Waters at the latter end of Pregnancy.
D The ſame contracted and grown thicker after the Waters are evacuated.
E.E The figure of the *Uterus* when *Pendulous*. In this Caſe if the *Membranes* break when the Patient is in an erect poſition, the Head of the *Fœtus* runs a risk of ſliding over and above the *Oſſa Pubis*, whence the Shoulders will be puſhed into the *Pelvis*.
F.F The figure of the *Uterus* when ſtretched higher than uſual, which generally occaſions Vomitings and Difficulty of Breathing. Conſult on this Subject Mr. *Levret* ſur le Mechaniſme de differentes Groſſeſſes.
G The *Os Pubis* of the left Side.
H.H The *Os Internum*.
I The *Vagina*.
K The left *Nympha*.
L The *Labium Pudendi* of the ſame Side.
M The remaining portion of the Bladder.
N The *Anus*.
O.P The left hip and thigh.

In this Period of Labour the *Os Uteri* being more and more ſtretched by the *Membranes* puſhing down, and beginning to extend the *Vagina*, a great quantity of Waters is forced down at the ſame time, and (if the *Membranes* break) are diſcharged; whence the *Uterus* contracts itſelf nearer to the Body of the *Fœtus*, which is here repreſented in a natural poſition with the *Vertex* reſting at the ſuperior part of the *Oſſa Pubis* and the forehead towards the right *Os Ilium*. As ſoon as the *Uterus* is in contact with the body of the *Fœtus*, the Head of the ſame is forced backwards towards the *Os Sacrum* from the line of the *Abdomen* B.G into that of the *Pelvis*, *viz.* from the uppermoſt F to near the end of the *Coccyx*, and is gradually puſhed lower as in the following Table.

If the *Membranes* do not break immediately upon their being puſhed into the *Vagina*, they ſhould be allowed to protrude ſtill further in order to dilate the *Os Externum*.

Vide Vol. I. Lib. I. Chap. 2. Sect. 2. Chap. 3. Sect. 3. Lib. III. Chap. 1. Sect. 1. 2. 4. Chap. 2. Sect. 3. Chap. 3. Sect. 4. No. 5. Vol. II. Coll. 10. No. 4. Caſe 3. 4. Coll. 14. Vol. III. Coll. 34. No. 2. Caſe 4.

B
A
F
C
E
D
F
C
E
H
H
G
M
K
L
N
P
O

A
G
G
F
B
B
H
H
I
I
E
E
D
D
C

THE

THIRTEENTH TABLE

In the ſame View and Section of the parts as in Table VI. ſhews the natural poſition of the Head of the *Fœtus* when ſunk down into the middle of the *Pelvis* after the *Os Internum* is fully opened, a large quantity of the Waters being protruded with the *Membranes* through the *Os Externum*, but prevented from being all diſcharged, from the Head's filling up the *Vagina*.

A The *Uterus* a little contracted, and thicker from ſome of the Waters being ſunk down before the Child, or diſcharged.
B.B The ſuperior parts of the *Oſſa Iliúm*.
C The inferior part of the *Rectum*.
D.D The *Vagina* largely ſtretched with the Head of the *Fœtus*.
E.E The *Os Internum* fully opened.
F A portion of the *Placenta*.
G.G The *Membranes*.
H.H The *Ligamenta lata*.
I.I The *Ligamenta rotunda*. Both theſe laſt ſtretched upwards with the *Uterus*.

The *Vertex* of the *Fœtus* being now down at the inferior part of the right *Os Iſchiúm*, and the wide part of the Head at the narrow and inferior part of the *Pelvis*, the forehead by the force of the pains is gradually moved backwards, and as it advances lower, the *Vertex* and *Occiput* turn out below the *Pubes*, as in the next Table. Hence may be learnt of what conſequence it is to know, that it is wider from ſide to ſide at the Brim of the *Pelvis*, than from the back to the forepart, and that it is wider from the fore to the hind Head of the Child, than from Ear to Ear.

Vide Vol. I. Lib. I. Chap. 1. Sect. 3. 5. alſo Lib. III. Chap. 3. Sect. 3. 4. No. 3. Vol. 2. Coll. 14.

THE

FOURTEENTH TABLE

In a ſimilar View and Section of the Parts with Table XII. ſhews the forehead of the *Fœtus* turned [in it's progreſſion downwards, from it's poſition in the former Table,] backwards to the *Os Sacrum*, and, the *Occiput* below the *Pubes*, by which means the narrow part of the Head is to the narrow part of the *Pelvis*, that is, between the inferior parts of the *Oſſa Iſchiûm*. Hence it may be obſerved, that though the diſtance between the inferior parts of the laſt mentioned Bones is much the ſame, as between the *Coccyx* and *Pubes*, yet as the Cavity of the *Pelvis* is much ſhallower at the anterior than lateral part, the *Occiput* of the *Fœtus*, when come down to the inferior part of either *Os Iſchiûm*, tûrns out below the *Pubes*; this anſwers the ſame end as if the *Pelvis* itſelf had been wider from the poſterior part than from ſide to ſide; the Head likewiſe enlarging the Cavity by forcing back the *Coccyx*, and puſhing out the external parts in form of a large Tumor, as is more fully deſcribed in the following Table.

Vide Vol. I. II. as referred to in the preceding Table.

A The *Uterus* contracted cloſely to the *Fœtus* after the Waters are evacuated.
B.C.D The *Vertebræ* of the Loins, *Os Sacrum* and *Coccyx*.
E The *Anus*.
F The left Hip.
G The *Perinæum*.
H The *Os Externum* beginning to dilate.
I The *Os Pubis* of the left ſide.
K The remaining portion of the Bladder.
L The poſterior part of the *Os Uteri*.

N. B. Although for the moſt part, at or before this period, the Waters are evacuated, yet it often happens, that more or leſs will be retained, and not all diſcharged, till after the delivery of the Child; occaſioned from the preſenting part of the *Fœtus* coming into cloſe contact with the lower or under part of the *Uterus*, *Vagina*, or *Os Externum*, immediately, or ſoon after the *Membranes* break.

B
A
C
I
K
L
D
H
G
E
F

TAB. XV.



THE

FIFTEENTH TABLE

Is intended principally to ſhew in what manner the *Perinæum*, and external parts are ſtretched by the Head of the *Fœtus*, in a firſt *pregnancy*, towards the end of Labour.

A The *Abdomen.*
B The *Labia Pudendi.*
C The *Clitoris*, and its *Præputium.*
D The hairy ſcalp of the *Fœtus* ſwelled at the *Vertex*, in a laborious caſe, and protruded to the *Os Externum.*
E.F The *Perinæum* and *Anus* puſhed out by the head of the *Fœtus* in form of a large *Tumor.*
G.G The parts that cover the *Tuberoſities* of the *Oſſa Iſchiûm.*
H The part that covers the *Os Coccygis.*

The *Perinæum* in this figure is ſtretched two inches, or double its length in the natural ſtate; but when the *Os Externum* is ſo much dilated by the head of the *Fœtus* as to allow the delivery of the ſame, the *Perinæum* is generally ſtretched to the length of three, and ſometimes four inches. The *Anus* is likewiſe lengthened an inch, the parts alſo between it and the *Coccyx* being much diſtended. All this ought to caution the young Practitioner never to precipitate the delivery at this time, but to wait, and allow the parts to dilate in a ſlow manner, as from the violence of the Labour-pains the ſudden delivery of the head of the *Fœtus* might endanger the laceration of the parts. The palm of the Operator's hand ought therefore to be preſſed againſt the *Perinæum*, that the head may be prevented from paſſing till the *Os Externum* is ſufficiently dilated, to allow its delivery without tearing the *Frœnum*, and parts betwixt that and the *Anus*, which are at this time very thin.

Vide Vol. I. Lib. III. Chap. 2. Sect. 2. Chap. 3. Sect. 4, No. 1. and Lib. 4. Chap. 1. Sect. 1. alſo Vol. II. Coll. 14. 24. Vol. III. Coll. 40.

THE

SIXTEENTH TABLE

And the three following ſhew in what Manner the Head of the *Fœtus* is helped along with the Forceps, as artificial Hands, when it is neceſſary to aſſiſt with the ſame for the ſafety of either Mother or Child. In this Table the Head is repreſented as forced down into the *Pelvis* by the Labour-pains, from its former Poſition in Table XII.

A.A.B.C The *Vertebræ* of the Loins, *Os Sacrum* and *Coccyx*.
D The *Os Pubis* of the left ſide.
E The remaining part of the Bladder.
F.F The *Inteſtinum Rectum*.
G.G.G The *Uterus*.
H The *Mons Veneris*.
I The *Clitoris* with the left *Nympha*.
X The *Corpus Cavernoſum Clitoridis*.
V The *Meatus Urinarius*.
K The left *Labium Pudendi*.
L The *Anus*.
N The *Perinæum*.
Q.P The left Hip and Thigh.
R The Skin and muſcular part of the Loins.

The Patient in this Caſe may be as in this Table, on her ſide with her Breech a little over the ſide or foot of the Bed, her Knees being likewiſe pulled up to her Belly, and a Pillow placed between them, care being taken at the ſame time that the parts are by a proper covering defended from the external air. If the hairy ſcalp of the *Fœtus* is ſo ſwelled that the ſituation of the Head cannot be diſtinguiſhed by the *Sutures* as in Table XXI. or if by introducing a Finger between the Head of the Child and the *Pubes*, or *Groins*, the Ear or back part of the Neck cannot be felt, the *Os Externum* muſt be gradually dilated in the time of the pains with the Operator's fingers, (previouſly lubricated with Hog's-lard) till the whole Hande an be introduced into the *Vagina*, and ſlipped up in a flattiſh form between the poſterior part of the *Pelvis* and Child's Head. This laſt is then to be raiſed up as high as is poſſible, to allow room for the Fingers to reach the Ear and poſterior part of the Neck. When the poſition of the Head is known, the Operator muſt withdraw his Hand, and wait to ſee if the ſtretching of the parts will renew or increaſe the Labour-pains, and allow more ſpace for the advancement of the Head in the *Pelvis*. If this however proves of no effect, the Fingers are again to be introduced as before, and one of the Blades of the Forceps (lubricated with lard) is then to be applied along the inſide of the Hand or Fingers, and left Ear of the Child, as repreſented in the Table. But if the *Pelvis* is diſtorted, and projects forward at the ſuperior part of the *Os Sacrum*, and the Forehead therefore cannot be moved a little backwards, in order to turn the Ear from that part of the *Pelvis* which prevents the End of the Forceps to paſs the ſame, in that caſe I ſay, the Blade muſt be introduced along the poſterior part of the Ear at the ſide of the diſtorted Bone. The Hand that was introduced is then to be withdrawn, and the Handle of the introduced Blade held with it as far back as the *Perinæum* will allow, whilſt the Fingers of the other Hand are introduced to the *Os Uteri*, at the *Pubes* or right *Groin*, and the other Blade placed exactly oppoſite to the former. This done, the Handles being taken hold of and joined together, the Head is to be pulled lower and lower every pain till the *Vertex*, as in this Table, is brought down to the inferior part of the left *Iſchiûm*, or below the ſame. The wide part of the Head being now advanced to the narrow part of the *Pelvis* betwixt the *Tuberoſities* of the *Oſſa Iſchiûm*, it is to be turned from the left *Iſchiûm* out below the *Pubes*, and the Forehead backwards to the concave part of the *Os Sacrum* and *Coccyx*, as in Table XVII. and afterwards the Head brought along and delivered as in Table XVIII. and XIX. But if it is found that the delivery will require a conſiderable degree of force from the Head's being large, or the *Pelvis* narrow, the Handles of the Forceps are to be tied together with a Fillet, as repreſented in this Table, to prevent their poſition being changed, whilſt the Woman is turned on her Back, as in Table XXIV. which is then more convenient for delivering the Head than when lying on the ſide.

This Table ſhews that the Handles of the Forceps ought to be held as far back as the *Os Externum* will allow, that the Blades may be in an imaginary line between that and the middle ſpace between the *Umbilicus* and the *Scrobiculus Cordis*. When the Forceps are applied along the Ears and Sides of the Head, they are nearer to one another, have a better hold, and mark leſs than when over the *Occipital* and *Frontal* Bones.

Vide Vol. I. Lib. III. Chap. 3. from Sect. 1. to 6. and Vol. II. Coll. 25, 26, 27. and 29.


R
A
A
G
F
G
B
F
C
L
N
G
E
H
D
X
I
V
K
P
Q


G
G
A
F
B
F
C
L
M
N
G
O
E
H
D
X
I
K
Q
R
S

THE

SEVENTEENTH TABLE

In the ſame View with the former repreſents in out-lines the Head of the *Fœtus* brought lower with the Forceps, and turned from the poſition in the former Table, in imitation of the natural progreſſion by the Labour-pains, which may likewiſe be ſuppoſed to have made this turn, before it was neceſſary to aſſiſt with the Forceps, this neceſſity at laſt ariſing from many of the cauſes mentioned in Vol. I.

In this View the poſition of the Forceps, along the Ears and narrow part of the Head, is more particularly expreſſed. It appears alſo that when the *Vertex* is turned from the left *Os Iſchiûm*, where it was cloſely confined, it is diſengaged by coming out below the *Pubes*, and the Forehead that was preſſed againſt the middle of the right *Os Iſchiûm*, is turned into the Concavity of the *Os Sacrum* and *Coccyx*. By this means the narrow part of the Head is now between the *Oſſa Iſchiûm*, or narrow part of the *Pelvis*, and as the *Occiput* comes out below the *Pubes*, the Head paſſes ſtill eaſier along. When the Head is advanced ſo low in the *Pelvis*, if the poſition cannot be diſtinguiſhed by the Sutures, it may for the moſt part be known by feeling for the back part of the neck of the *Fœtus*, with a Finger introduced betwixt the *Occiput* and *Pubes*, or towards one of the *Groins*. If the Head is ſqueezed into a longiſh form, as in Table XXI. and has been detained many Hours in this poſition, the pains not being ſufficient to compleat the Delivery, the Aſſiſtance of the Forceps muſt be taken to ſave the Child, though the Woman may be in no danger. But if the Head is high up in the *Pelvis*, as in the former Table, the Forceps ought not to be uſed except in the moſt urgent neceſſity.

This Table alſo ſhews that the handles of the Forceps are ſtill to be kept back to the *Perinæum*, and when in this poſition are in a line with the upper part of the *Sacrum*, and if held more backwards, when the Head is a little higher, would be in a line with the *Scrobiculus Cordis*. If the Forceps are applied when the Head is in this poſition, they are more eaſily introduced when the Patient is in a ſupine poſition, as in Table XXIV. Neither is it neceſſary to tie the handles, which is only done to prevent their alteration when turning the Woman from her ſide to her back.

As I have had ſeveral Caſes where a longer ſort of Forceps that are curved upwards are of great uſe to help along the Head, when the Body is delivered firſt, as in Table XXXV. the ſame are repreſented here by dotted lines. They may be uſed in laborious Caſes as well as the others, but are not managed with the ſame eaſe.

Moſt of the parts of this Table being marked with the ſame letters as the former, the deſcriptions there given will anſwer in this, except the following.

L.M The *Anus*.

M.N The *Perinæum*.

O The common *Integuments* of the *Abdomen*.

R The ſhort Forceps.

S The long curved Forceps, the firſt of theſe is eleven Inches long, and the laſt twelve Inches and a half, which I have after ſeveral alterations found ſufficient, but this need not confine others who may chooſe to alter them from this ſtandard. *Vide* Tab. XXXVII.

THE

EIGHTEENTH TABLE

In the ſame View and Section of the parts, ſhews the Head of the *Fœtus* in the ſame poſition, but brought lower down with the Forceps than in the former Table; for in this the *Os Externum* is more open, the *Occiput* come lower down from below the *Pubes*, and the Forehead paſt the *Coccyx*, by which both the *Anus* and *Perinæum* are ſtretched out in form of a large Tumor, as in Table XV.

When the Head is ſo far advanced, the Operator ought to extract with great Caution, leſt the parts ſhould be torn. If the Labour-pains are ſufficient, the Forehead may be kept down, and helped along in a ſlow manner by preſſing againſt it with the Fingers on the external parts below the *Coccyx*: at the ſame time the Forceps being taken off, the Head may be allowed to ſtretch the *Os Externum* more and more in a gradual manner, from the force of the Labour-pains, as well as aſſiſtance of the Fingers. But if the former are weak and inſufficient, the aſſiſtance of the Forceps muſt be continued. *Vide* the deſcription of the parts in Table XVI. S.T in this repreſent the left ſide of the *Os Uteri*. The dotted lines demonſtrate the ſituation of the Bones of the *Pelvis* on the right ſide, and may ſerve as an Example for all the lateral Views of the ſame.

a.b.c.h. The out-lines of the *Os Ilium*.
D.e.f. The ſame of the *Pubis* and *Iſchium*.
i.i.k. The *Acetabulum*. And
m.n. The *Foramen Magnum*.

Vide Vol. I. Lib. III. Chap. 5. Sect. 3. Vol. II. Collect. 25.


a
G
R
A
G
F
b
h
G
i
S
T
G
B
F
c
E
f
k
l
m
H
D
P
I
K
L
e
M
Q
N

R
B
Q
C
F
L
M
N
S
T
G
F
A
D
E
K
H
P
G
G
O

THE

NINETEENTH TABLE

In the ſame View and Section of the *Pelvis*, is intended by out-lines to ſhew that as the external parts are ſtretched, and the *Os Externum* is dilated, the *Occiput* of the *Fœtus* riſes up with a ſemicircular turn from out below the *Pubes*, the under part of which Bones are as an Axis, or Fulcrum, on which the back part of the Neck turns, whilſt at the ſame time, the Forehead and Face in their turn upwards, diſtend largely the parts between the *Coccyx*, and *Os Externum*. This is the method obſerved by Nature, in ſtretching theſe parts in Labour, and as Nature is always to be imitated, the ſame method ought to be followed, when it is neceſſary to help along the Head with the Forceps.

Vide the three former Tables for the Deſcriptions, and References.

THE

TWENTIETH TABLE

In the ſame Section of the parts, but with a view of the right ſide; ſhews the Head of the *Foetus* in the contrary poſition to the three laſt figures, the *Vertex* being here in the Concavity of the *Sacrum*, and the Forehead turned to the *Pubes*.

A.B The *Vertebræ* of the Loins, *Os Sacrum*, and *Coccyx*.
C The *Os Pubis* of the right ſide.
D The *Anus*.
E The *Os Externum* not yet begun to ſtretch.
F The *Nympha*.
G The *Labium Pudendi* of the right ſide.
H The Hip and Thigh.
I.I The *Uterus* contracted the Waters being all diſcharged.

When the Head is ſmall, and the *Pelvis* large, the *Parietal* Bones and the Forehead will, in this Caſe, as they are forced downwards by the Labour-pains, gradually dilate the *Os Externum*, and ſtretch the parts between that and the *Coccyx* in form of a large Tumor, as in Table XV. till the Face comes down below the *Pubes*, when the Head will be ſafely delivered. But if the ſame be large and the *Pelvis* narrow, the difficulty will be greater, and the Child in danger; as in the following Table.

Vide Vol. I. Lib. III. Chap. 3. Sect. 4. No. 3. Vol. II. Col. 16. No. 2.

I
A
I
C
G
F
E
D
B
H


N
M
O
K
L

THE

TWENTY-FIRST TABLE

Shews the Head of the *Fœtus* in the ſame poſition as in the former Table, but being much larger, it is by ſtrong Labour-pains ſqueezed into a longiſh form with a Tumor on the *Vertex*, from the long compreſſion of the Head in the *Pelvis*. If the Child cannot be delivered with the Labour-pains, or turned and brought footling, the Forceps are to be applied on the Head as deſcribed in this figure, and brought along as it preſents; but if that cannot be done without running the riſque of tearing the *Perinæum*, and even the *Vagina* and *Rectum* of the Woman, the Forehead muſt be turned backwards to the *Sacrum*. To do this more effectually, the Operator muſt graſp firmly with both Hands the handles of the Forceps, and at the ſame time puſhing upwards raiſe the Head as high as poſſible, in order to turn the Forehead to one ſide, by which it is brought into the natural poſition; this done the Head may be brought down and delivered as in Table XVI. &c.

Vide Vol. I Lib. III. Chap. 3. Sect. 4. No. 2. and Voll. II. Coll. 28. alſo the former Table for the Deſcription of the parts, except

K The Tumor on the *Vertex*. The ſame compreſſion and elongation of the Head as well as the Tumor on the *Vertex*, may be ſuppoſed to happen in a greater or leſs degree in the XVI. XVII. XVIII. XIX. Tables as well as in this, where the Difficulty proceeds from the Head being large, or the *Pelvis* narrow. *Vide* Table XXVII. and XXVIII.

L The Forceps. Sometimes the Forehead may be moved to the natural poſition by the aſſiſtance of the Fingers, or only one Blade of the Forceps. The Forceps may either be the ſtraight kind, or ſuch as are curved to one ſide, when it is neceſſary to uſe one or both Blades.

M The *Veſica Urinaria* much diſtended with a large quantity of *Urine* from the long preſſure of the Head againſt the *Urethra*, which ſhews that the *Urine* ought to be drawn off with a *Catheter* in ſuch extraordinary Caſes before you apply the Forceps, or in præternatural Caſes, where the Child is brought footling.

N The under part of the *Uterus*.

O.O The *Os Uteri*.

THE

TWENTY-SECOND TABLE

Shews in a front View of the parts, the Forehead of the *Eætus* presenting at the Brim of the *Pelvis*, the Face being turned to one side, the *Fontanelle* to the other, and the Feet and Breech stretched towards the *Fundus Uteri*.

A.A The superior part of the *Ossa Ilium*.
B The *Anus*.
C The *Perinæum*.
D The *Os Externum*, the thickness of the posterior part before it is stretched with the Head of the Child.
E.E.E The *Vagina*.
F The *Os Uteri* not yet fully dilated.
G.G.G The *Uterus*.
H The *Membrana Adiposa*.

If the Face is not forced down, the Head will sometimes come along in this Manner, in which Case the *Vertex* will be flattened, and the Forehead raised in a conical Form, and when the Head comes down to the lower part of the *Pelvis*, the Face or *Occiput* will be turned from the side, and come out below the *Pubes*. But if the Head is large and cannot be delivered by the pains, or if the wrong position cannot be altered, the Child must, if possible, be brought footling, or delivered with the Forceps.

Vide Vol. I. Lib. III. Chap. 2. Sect. 3. Chap. 3. Sect. 4. No. 3. Vol. II. Coll. 16. No. 4. Coll. 28.

G
A
A
G
G
F
E
H
D
C
B

A
G
G
H
C
F
B
E
D

THE

TWENTY-THIRD TABLE

Shews in a lateral View the Face of the Child presenting, and forced down into the lower part of the *Pelvis*, the Chin being below the *Pubes*, and the *Vertex* in the Concavity of the *Os Sacrum*: the Waters likewise being all discharged, the *Uterus* appears closely joined to the Body of the Child, round the Neck of which is one circumvolution of the *Funis*.

A.B The *Vertebræ* of the Loins, *Os Sacrum*, and *Coccyx*.
C The *Os Pubis* of the left side.
D The inferior part of the *Rectum*.
E The *Perinæum*.
F The left *Labium Pudendi*.
G.G.G The *Uterus*.

When the *Pelvis* is large, the Head, if small, will come along in this position, and the Child be saved: for as the Head advances lower, the Face and Forehead will stretch the parts between the *Frænum Labiorum* and *Coccyx* in form of a large Tumor. As the *Os Externum* likewise is dilated, the Face will be forced through it, the under part of the Chin will rise upwards over the anterior part of the *Pubes*, and the Forehead, *Vertex*, and *Occiput*, turn up from the parts below. If the Head however is large, it will be detained either when higher or in this position. In this case, if the position cannot be altered to the natural, the Child ought to be turned, and delivered footling. If the *Pelvis* however is narrow, and the Waters not all gone, the *Vertex* should, if possible, be brought to present; but if the *Uterus* is so closely contracted, that this cannot be effected, on account of the strong pressure of the same, and slipperiness of the Child's Head, in this case the Method directed in the following Table is to be taken.

THE

TWENTY-FOURTH TABLE

Repreſents in the lateral View, the Head of the *Fœtus* in the ſame poſition as in the former Table; but the delivery is ſuppoſed to be retarded from the largeneſs of the Head, or a narrow *Pelvis*.

In this Caſe, if the Head cannot be raiſed, and puſhed up into the *Uterus*, it ought to be delivered with the Forceps, in order to ſave the Child. This poſition of the Chin to the *Pubes* is one of the ſafeſt caſes, where the Face preſents, and is moſt eaſily delivered with the Forceps, the manner of introducing of which over the Ears is ſhewn in this Table. The Patient muſt lie on her Back with her Breech a little over the Bed, her Legs and Thighs being ſupported by an aſſiſtant ſitting on each ſide. After the parts have been ſlowly dilated with the Hand of the Operator, and the Forceps introduced, and properly fixed along the Ears of the Child, the Head is to be brought down by degrees, that the parts below the *Os Externum* may be gradually ſtretched: the Chin then is to be raiſed up over the *Pubes*, whilſt the Forehead, *Fontanelle*, and *Occiput*, are brought out ſlowly from the *Perinæum* and *Fundament* to prevent the ſame from being hurt, or lacerated. But if the *Fœtus* can neither be turned, nor extracted with the Forceps, the delivery muſt be left to the Labour-pains, as long as the Patient is in no danger; but if danger is apparent, the Head muſt be delivered with the curved Crotchets. *Vide* Table XXXIX.

When the Face preſents, and the Chin is to the ſide of the *Pelvis*, the Patient muſt lie on her ſide, and after the Forceps are fixed along the Ears, the Chin is to be brought down to the lower part of the *Os Iſchiûm*, and then turned out below the *Pubes*, and delivered in a ſlow manner as above.

Vide Vol. II. Coll. 16. No. 6. as alſo Tables XVI. XVII. XVIII. and XIX. for the Deſcription of the parts.



A
B

THE

TWENTY-FIFTH TABLE

Shews in a lateral View of the right ſide, the Face of the *Fœtus* preſenting, as in Table XXIII. but in the contrary poſition, that is, with the Chin to the *Os Sacrum*, and the *Bregma* to the *Pubes*, the Waters evacuated, and the *Uterus* contracted.

A The *Os Externum* not yet begun to ſtretch.
B The *Anus*. *Vide* Table XX. for the further Deſcription of the parts.

In ſuch caſes, as well as in thoſe of the laſt mentioned Table, if the Child is ſmall, the Head will be puſhed lower with the Labour-pains, and gradually ſtretch the lower part of the *Vagina*, and the external parts, by which means, the *Os Externum* will be more and more dilated, 'till the *Vertex* comes out below the *Pubes*, and riſes up on the outſide; in which caſe the delivery is then the ſame, as in natural Labours. But if the Head is large, it will paſs along with great Difficulty, whence the Brain, and Veſſels of the Neck, will be ſo much compreſſed and obſtructed, as to deſtroy the Child. To prevent which, if called in time, before the Head is far advanced in the *Pelvis*, the Child ought to be turned, and brought footling. If the Head however is low down, and cannot be turned, the delivery is then to be performed with the Forceps, either by bringing along the Head as it preſents, or as in the following Table. See the References in the preceding Table.

THE

TWENTY-SIXTH TABLE

Repreſents by out-lines in a lateral View of the left ſide of the Subject, the *Fœtus* in the ſame Situation as in the former Table.

The Head here is ſqueezed into a very oblong form, and though forced down ſo as fully to dilate the *Os Externum*, yet the *Vertex* and *Occiput* cannot be brought ſo far down, as to turn out from below the *Pubes*, (as in the foregoing Table) without tearing the *Perinæum* and *Anus*, as well as the *Vagina* and *Rectum*.

The beſt method in this caſe, after either the ſhort or long curved Forceps have been applied along the Ears, (as repreſented in the Table) is to puſh the Head as high up in the *Pelvis* as is poſſible, after which the Chin is to be turned from the *Os Sacrum* to either *Os Iſchiûm*, and afterwards brought down to the inferior part of the laſt mentioned Bone. This done, the Operator muſt pull the Forceps with one Hand, whilſt two Fingers of the other are fixed on the lower part of the Chin or Under-jaw, to keep the Face in the middle, and prevent the Chin from being detained at the *Os Iſchiûm*, as it comes along, and in this manner move the Chin round with the Forceps, and the above Fingers, till brought under the *Pubes*, which done the Head will be eaſily extracted, as in Table XXIV.

If, before aſſiſtance has been called, the Head is ſo ſqueezed down into the *Pelvis*, that it is impoſſible to move the Chin from the *Sacrum* to either *Os Iſchiûm*, ſo as to deliver with the Forceps for the Safety of the Child, the Operator muſt wait with Patience, as long as the Woman is not in danger, or there is no certainty of the death of the *Fœtus:* But if the Patient runs the leaſt Riſque, the Head muſt be delivered with the Crotchet.

In general, with reſpect to the poſture of the Woman in the application of the Forceps, when the Ears are to the ſides of the *Pelvis*, the Forceps, as was obſerved in Table XXIV. are moſt eaſily introduced when the Patient lies upon her Back, and her Breech over the ſide of the Bed; but when the Ear is to the *Pubes* or *Groin*, they are better applied when the Patient lies on her ſide, as was obſerved in the caſes where the *Vertex* preſented.

Vide Table XXIV. for the Deſcription of the Parts and the References, alſo Table XXXIX. for the manner of uſing the Crotchet.


A
B
C
D
E

THE

TWENTY-SEVENTH TABLE

Gives a lateral internal View of a diſtorted *Pelvis*, divided longitudinally, with the Head of a *Fœtus* of the ſeventh Month paſſing the ſame. *Vide* the Explanation of Table III.

A.B.C The *Os Sacrum* and *Coccyx*.
D The *Os Pubis* of the left ſide.
E The *Tuberoſity* of the *Os Iſchiûm*, of the ſame ſide.

The Head of the *Fœtus* here, though ſmall, is with Difficulty ſqueezed down into the *Pelvis*, and changed from a round to an oblong Form, before it can paſs, there being only the Space of two Inches and one quarter between the projection of the ſuperior part of the *Sacrum*, and the *Oſſa Pubis*. If the Head is ſoon delivered, the Child may be born alive: but if it continues in this manner many Hours, it is in danger of being loſt, on account of the long Preſſure on the Brain. To prevent which, if the Labour-pains are not ſufficiently ſtrong, the Head may be helped along with the Forceps, as directed in Table XVI.

This figure may ſerve as an example of the extreme degree of diſtortion of the *Pelvis*, between which, and the well-formed one, are many intermediate degrees, according to which the Difficulty of Delivery muſt increaſe, or diminiſh, as well as from the diſproportion of the *Pelvis* and Head of the *Fœtus*; all which Caſes require the greateſt Caution, both as to the Management and Safety of the Mother and Child.

Vide Vol. I. Lib. III. Chap. 2. Sect. 3. No. 5. Chap. 3. Sect. 4. No. 3. Vol. II. Coll. 21. No. 1. and Coll. 29.

THE

TWENTY-EIGHTH TABLE

Gives a ſide View of a diſtorted *Pelvis*, as in the former Table, with the Head of a full grown *Foetus*, ſqueezed into the Brim, the *Parietal* Bones decuſſating each other, and compreſſed into a conical Form.

A.B.C The *Os Sacrum* and *Coccyx*.
D The *Os Pubis* of the left ſide.
E The *Tuberoſity* of the *Os Iſchiûm*.
F The *Proceſſus Acutus*.
G The *Foramen Magnum*.

This Table ſhews the impoſſibility in ſuch a caſe to ſave the Child, unleſs by the *Cæſarian* Operation, which however ought never to be performed, excepting when it is impracticable to deliver at all by any other method. Even in this caſe, after the upper part of the Head is diminiſhed in bulk, and the Bones are extracted, the greateſt Force muſt be applied in order to extract the Bones of the Face, and Baſis of the Skull, as well as the Body of the *Fœtus*.

Vide Vol. I. Lib. III. Chap. 3. Sect. 7. Chap. 5. Sect. 3. and Vol. III. Coll. 31. 39.


A
B
C
D
E
F
G



THE

TWENTY-NINTH TABLE

Repreſents in a front View of the *Pelvis*, as in Table XXII. the Breech of the *Fœtus* preſenting, and dilating the *Os Internum*, the *Membranes* being too ſoon broke. The fore-parts of the Child are to the poſterior part of the *Uterus*, and the *Funis* with a knot upon it ſurrounds the Neck, Arm, and Body.

Some time after this and the following Tables were engraved, Dr. *Kelly* ſhewed me a Subject he had opened, where the Breech preſented itſelf, and lay much in the ſame poſition with its Body as in the Ninth Table, ſuppoſing the Breech in that Figure, turned down to the *Pelvis*, and the Head up to the *Fundus Uteri*.

I have ſometimes felt, in theſe Caſes, [when Labour was begun, and before the Breech was advanced into the *Pelvis*,] one Hip at the *Sacrum*, the other reſting above the *Os Pubis*, and the private Parts to one Side; but before they could advance lower, the *Nates* were turned to the ſides and wide part of the Brim of the *Pelvis*, with the private parts to the *Sacrum*, as in this Table; though ſometimes to the *Pubis*, as in the following Table. As ſoon as the Breech advances to the lower part of the Baſon, the Hips again return to their former poſition, *viz.* one Hip turned out below the *Os Pubis*, and the other at the back Parts of the *Os Externum*.

N.B. In this caſe the Child, if not very large, or the *Pelvis* narrow, may be often delivered alive by the Labour-pains; but if long detained at the inferior part of the *Pelvis*, the long preſſure of the *Funis* may obſtruct the circulation. In moſt Caſes where the Breech preſents, the effect of the Labour-pains ought to be waited for, 'till at leaſt they have fully dilated the *Os Internum* and *Vagina*, if the ſame have not been ſtretched before with the Waters and *Membranes*. In the mean time, whilſt the Breech advances, the *Os Externum* may be dilated gently during every Pain, to allow room for introducing a Finger or two of each Hand to the outſide of each Groin of the *Fœtus*, in order to aſſiſt the Delivery when the *Nates* are advanced to the lower part of the *Vagina*. But if the *Fœtus* is larger than uſual, or the *Pelvis* narrow, and after a long time, and many repeated pains, the Breech is not forced down into the *Pelvis*, the Patient's Strength at the ſame time failing, the Operator muſt in a gradual manner open the Parts, and having introduced a Hand into the *Vagina*, raiſe or puſh up the Breech of the *Fœtus*, and bring down the Legs and Thighs. If the *Uterus* is ſo ſtrongly contracted that the Legs cannot be got down, the largeſt end of the blunt Hook is to be introduced, as directed in Table XXXVII. As ſoon as the Breech or Legs are brought down, the Body and Head are to be delivered as deſcribed in the next Table, only there is no Neceſſity here to alter the poſition of the Child's Body.

Vide Vol. I. Lib. III. Chap. 4. Sect. 1, 2. Vol. III. Coll. 32.

The deſcription of the Parts in this, and the following Table, is the ſame as in Table XXII. only the dotted Lines in this deſcribe the place of the *Oſſa Pubis*, and anterior parts of the *Oſſa Iſchiûm* which are removed, and may ſerve in this reſpect, as an example for all the other front Views, where, without disfiguring the Table, they could not be ſo well put in.

THE

THIRTIETH TABLE

Shews in the ſame View, and with the ſame References as the former, the Breech of the *Fœtus* preſenting, with this Difference however, that the fore-parts of the Child are to the fore-part of the *Uterus*. In this Caſe when the Breech coming double as it preſents is brought down to the Hams, the Legs muſt be extracted, a Cloth wrapped round them, and the fore-parts of the Child turned to the back-parts of the Woman. If a Pain ſhould in the mean time force down the Body of the Child, it ought to be puſhed up again in turning, as it turns eaſier when the Belly is in the *Pelvis*, than when the Breaſt and Shoulders are engaged, and as ſometimes the Face and Forehead are rather towards one of the Groins, a quarter turn more brings theſe parts to the ſide of the *Pelvis*, and a little backwards, after which the Body is to be brought down. If the Child is not large, the Arms need not be brought down, and the Head may be delivered by preſſing back the Shoulders and Body of the Child to the *Perinæum*, and whilſt the Chin and Face are within the *Vagina*, to bring the *Occiput* out from below the *Pubes*, according to *Daventer*'s method. Or the Operator may introduce a Finger or two into the Mouth, or on each ſide of the Noſe, and ſupporting the Body on the ſame Arm, fix two Fingers of the other Hand over the Shoulders, on each ſide of the Child's Neck, and in this manner raiſe the Body over the *Pubes*, and bring the Face and Forehead out with a ſemicircular turn upwards, from the under part of the *Os Externum*. All this may be eaſily done when the Woman lies on her Side; but if the Child is large, and the *Pelvis* narrow, it is better to turn the Patient on her Back, as deſcribed in Table XXIV. and after the Legs and Body are extracted as far as the Shoulders, the Arms are to be cautiouſly brought down, and the Head delivered. If the Woman has ſtrong Pains, and when by the felt Pulſation of the Veſſels of the *Funis Umbilicalis*, or the ſtruggling Motions of the *Fœtus*, it is certain that the Child is ſtill alive, wait with Patience for the Aſſiſtance of the Labour; but if that and the Hand are inſufficient, and the Pulſation of the *Funis* turns weaker, and if the Child cannot be brought double, the Breech muſt be puſhed up, and if the Reſiſtance of the *Uterus* is ſo great, as to prevent the Extraction of the Legs, the Patient ought to be turned on her Knees and Elbows. When the Legs are thus brought down, the Woman, if needful, is to be again turned to her Back, to allow more Freedom to deliver the Body and Head, as before deſcribed. If the Head after ſeveral Trials cannot be delivered, without endangering the Child, from overſtraining the Neck, the long curved Forceps ought to be applied, as in Table XXXV. If theſe fail, and the Patient is not in Danger, ſome Time may be allowed for the Effect of the Labour-pains, which likewiſe proving inſufficient, the Crotchet muſt be uſed as in Table XXXIX. and when it is certain that the Child is dead, or that there is no Poſſibility of ſaving it.

B
A
F
A
F
B
E
O
C
C
F
E
O
E
O
E
D

THE

THIRTY-FIRST TABLE

Repreſents, in a front View of the *Pelvis*, the *Fœtus* compreſſed by the Contraction of the *Uterus* into a round form, the fore-parts of the former being towards the inferior part of the latter, and one Foot and Hand fallen down into the *Vagina*. In this figure the anterior part of the *Pelvis* is removed by a longitudinal Section through the middle of the *Foramen Magnum*.

A.A The ſuperior parts of the *Oſſa Ilium*.
B B The *Uterus*.
C The Mouth of the *Womb* ſtretched, and appearing in
O.O.O.O The *Vagina*.
D The inferior and poſterior part of the *Os Externum*.
E.E.E.E The remaining parts of the *Oſſa Pubis* and *Iſchium*.
F.F.F.F The *Membrana Adipoſa*.

This and the three following Tables repreſenting four different præternatural poſitions of the *Fœtus* in *Utero*, may ſerve as Examples for the manner of Delivery in theſe as well as in all other præternatural Caſes.

In all præternatural Caſes, the *Fœtus* may be eaſily turned and delivered by the Feet, if known before the *Membranes* are broke, and the Waters diſcharged; or if the *Pelvis* is narrow, and the Patient is ſtrong, the Head, if large, may be brought down ſo as to preſent in the natural Way; but if all the Waters are diſcharged, and the *Uterus* is ſtrongly contracted to the Body of the *Fœtus*, this laſt Method can ſeldom take place, on account of the ſtrong Preſſure of the *Uterus*, and Slipperineſs of the Child's Head.

In the preſent Caſe, the Woman may either be laid on her Back or Side, as deſcribed in Table XVI. and XXIV. and the Operator having ſlowly dilated the *Os Externum* with his Fingers, muſt introduce the ſame into the *Vagina*, and puſh up into the *Uterus* the parts of the *Fœtus* that preſent; or if there is ſpace for it, his Hand may paſs in order to dilate the *Os Internum* if not ſufficiently ſtretched previouſly by the *Membranes*, and Waters. This done he muſt advance his Hand into the *Uterus*, to know the poſition of the *Fœtus*, and as the Breech is rather lower than the Head, ſearch for the other Leg, and bring down both Feet without the *Os Externum*. A Cloth muſt then be wrapped round them, and having graſped them with one Hand, he is to introduce the other into the *Uterus*, in order to raiſe the Head of the *Fœtus*, whilſt the Legs and Thighs are pulled down by the Hand that holds the Feet. When the Head is raiſed, and does not fall down again, the Hand of the Operator may be withdrawn from the *Uterus*, and the Delivery completed, as directed in the two former Tables. By the artleſs method of taking hold and pulling one or both Feet, the Breech may come down and the Head riſe to the *Fundus*, but if this ſhould not happen, there will be great danger of overſtraining the *Fœtus*, which is prevented by the former method. If the *Membranes* are broken, before the *Os Uteri* is largely opened, and the Hand of the Operator cannot be introduced, which ſometimes happens in a firſt Pregnancy, the parts of the *Fœtus* ſhould be allowed to protrude ſtill further, by which means the rigidity of the *Os Internum* will in time be leſſened.

Vide Vol. I. and III. on præternatural Labours.

THE

THIRTY-SECOND TABLE

Repreſents in the ſame View with the former, the *Fœtus* in the contrary Poſition; the Breech and Fore-parts being towards the *Fundus Uteri*, the left Arm in the *Vagina*, and fore Arm without the *Os Externum*, the Shoulder being likewiſe forced into the *Os Uteri*.

The Operator in this Caſe muſt introduce his Fingers between the back part of the *Vagina* and the Arm of the *Fœtus*, in order to raiſe the Shoulder and make room for paſſing his Hand into the *Uterus* to diſtinguiſh the Poſition. This being known, he ought to puſh up the Shoulder to that part of the *Uterus*, where the Head is lodged, in order to raiſe the ſame to the *Fundus*. If the Body of the *Fœtus* does not move round, and thereby lie in a more convenient Poſition for bringing down the Legs, the Hand of the Operator ought to be puſhed up ſtill higher to ſearch for, and take hold of the Feet, which are to be brought down as far as is poſſible. If this ſhould not change the Poſition, the Shoulder is to be puſhed up, and the Legs pulled down alternately, 'till they are brought down into the *Vagina*, or without the *Os Externum*, after which the Delivery may be completed as in the former Caſe.

If the Feet cannot be brought down lower than into the *Vagina*, a Nooſe may be introduced over both Ankles, by which the Legs are brought lower by pulling the Nooſe with one Hand, whilſt the other, previouſly introduced into the *Uterus*, puſhes up the Shoulders and Head. By this double force the Poſition of the *Fœtus* is to be altered, and the Delivery effected. In theſe Caſes, as the Shoulder is raiſed to the *Fundus*, the Arm commonly returns into the *Uterus*; but if the Arm is ſo ſwelled as to prevent the introduction of the Operator's Hand, and cannot be folded up or returned into the *Uterus*, it muſt be taken off at the Shoulder, or Elbow, in order to deliver and ſave the Woman. If both Arms come down, when the Breaſt preſents, the Methods above deſcribed are to be uſed.

Vide the Explanations and References of the foregoing Table to illuſtrate this and the following.

A
B

THE

THIRTY-THIRD TABLE

Exhibits in the ſame View likewiſe of the *Pelvis* with the former, a third Poſition of the *Fœtus* when compreſſed into the round Form, the Belly, *viz.* or *Umbilical* Region, preſenting at the *Os Internum*, and the *Funis* fallen down into the *Vagina*, and appearing at the *Os Externum*.

The Delivery in this Caſe is to be effected as in the former Table, by puſhing up the Breaſt, and bringing down the Legs. When the Belly preſents, it is eaſier coming at the Legs, than when the Breaſt preſents, becauſe in the former Caſe the Head is nearer to the *Fundus Uteri*, and the Legs and Thighs lower. If the Belly or Breaſt is forced down into the lower part of the *Pelvis*, the Child will be in danger from the bending of the *Vertebræ*, and the Preſſure of the ſpinal Marrow, ſo great Force is alſo required to raiſe theſe Parts up into the *Uterus*, in order to come at the Feet, that it will ſometimes be neceſſary to turn the Woman to her Knees and Elbows, to diminiſh the Reſiſtance of the *Abdominal* Muſcles. When the *Funis* comes down without the *Os Externum*, if there is a Pulſation felt, it muſt immediately be replaced, and kept warm in the *Vagina*, to preſerve the Circulation, and prevent a Stagnation, from its being expoſed to the cold Air. If the *Funis* comes down when the Head preſents, the Child is in danger if not ſpeedily delivered with the Pains, or brought Footling.

See the two former Tables for the Explanations, and References.

THE

THIRTY-FOURTH TABLE

Shews, in a lateral View of the *Pelvis*, one of the moſt difficult præternatural Caſes. The left Shoulder Breaſt and Neck of the *Fœtus* preſenting, the Head reflected over the *Pubes* to the right Shoulder and Back, and the Feet and Breech ſtretched up to the *Fundus*, the *Uterus* contracted at the ſame time, in form of a long Sheath, round the Body of the *Fœtus*.

A.B.C The *Os Sacrum* and *Coccyx*.
D The *Os Pubis* of the left Side.
E Part of the *Urinary* Bladder.
F The *Rectum*.
H.I.K The Private Parts.
M The *Anus*.
M.N The *Perinæum*.
V The *Meatus Urinarius*.
O The *Os Uteri* not yet opened, and ſituated backwards towards the *Rectum* and *Coccyx*.
R.S The ſame repreſented in dotted Lines, as opened when the Labour is begun.
T.U The ſame more fully dilated, but nearer the poſterior than anterior part of the *Pelvis*.
W.P The ſame not fully ſtretched at the Fore-part, tho' intirely obliterated at the Back-part, the *Uterus* and *Vagina* being there only ſometimes one continued Surface.

Hence it appears why the anterior part of the *Os Uteri* is frequently protruded before the Head of the *Fœtus* at the *Pubes*, which, if it retards Delivery, is removed by ſliding it up with a Finger or two between the Head and laſt mentioned Part. *Vide* Table IX. X. XI. XII. XIII.

The manner of Delivery in the Poſition of the *Fœtus*, as repreſented in this Table, is to endeavour with the Hand to force up the part preſenting, in order to raiſe the Head to the *Fundus*. If this is impoſſible from the ſtrong Contraction of the *Uterus*, the Operator muſt puſh up his Hand in a ſlow and cautious manner along the Breaſt and Belly of the Child, in order to come at the Legs and Feet, which are to be taken hold of, and brought down as far as the Poſition of the *Fœtus* will admit of. The Body is then to be moved round by puſhing up the lower Parts, and pulling down the upper, till the Feet are brought without the *Os Externum*, and the Delivery completed as in Table XXXI. But if the Feet cannot be got down, ſo as to be taken hold of without the *Os Externum*, a Nooſe muſt be fixed over the Ankles, as in Table XXXII.

Vide Vol. I. and III. as directed in Table XXXI.


A
B
C
D
E
F
H
I
K
M
N
O
P
Q
R
S
T
U
V
W

D
F
E
A
F
E
F
B
H
G
I
C
C

THE

THIRTY-FIFTH TABLE

Shews, in a lateral View of the *Pelvis*, the Method of aſſiſting the Delivery of the Head of the *Fœtus* with the long curved Forceps, in præternatural Caſes, when it cannot be done with the Hands, as deſcribed in Table XXIX. and XXX.

A The three loweſt *Vertebræ* of the Loins, with the *Os Sacrum* and *Coccyx*.
B The *Os Pubis* of the left Side.
C.C The *Perinæum* and *Anus* preſſed backwards with the Forceps.
D The *Inteſtines*.
E.E The *Parietes* of the *Abdomen*.
F.F.F The *Uterus*.
G the poſterior part of the *Os Uteri*.
H The *Rectum*.
I The *Vagina*.

After the Body and Arms of the Child are delivered, and the different Methods uſed to bring down the Head with the Hands, as directed in the above Table, and more fully deſcribed in Vol. I. and III. the following Method is to be tried in order to ſave the Child, which muſt otherwiſe be loſt by overſtraining the Neck and ſpinal Marrow. The Woman being in the ſupine Poſition, as in Table XXIV. one of the aſſiſtants ought to hold the Body and Arms of the Child up towards the *Abdomen* of the Woman, to give more room to the Operator, who having introduced one hand up to the Child's Face, and moved it from the Side a little backwards, for the eaſier Application of the Forceps along the Sides of the Head, muſt then turn his hand to one of the Ears, and introduce one of the Blades, with the other hand between the ſame and the Head, with the curved Side towards the *Pubes*, as in this Table. This done, the hand is to be brought down to hold the Handle of the Blade of the Forceps, 'till the other Hand is introduced to the other ſide of the Head, by which means the ſame is preſſed againſt the Blade that is up, and which is thus prevented from ſlipping, whilſt the other Hand introduces the ſecond Blade on the oppoſite Side. The Blades being thus introduced, care muſt be taken, that in joining them no part of the *Vagina* is locked in. After the Forceps are firmly fixed along the Sides of the Head, the Face and Forehead muſt be turned again to the ſide of the Brim of the *Pelvis*, by which means the wide part of the Head is to the wide part of the Brim. This done the Head is to be brought lower, and the Force gradually increaſed, according to the Reſiſtance from the largeneſs of the Head, or narrowneſs of the *Pelvis*. The Forehead, when brought low enough down, is then to be turned into the Concavity of the *Os Sacrum* and *Coccyx*, the Handles of the Forceps raiſed upwards, and the ſame Caution uſed in bringing the Head through the *Os Externum*, as deſcribed in Table XIX. and XXX. By this Method the Head will be delivered, the Child frequently ſaved, and the uſe of the Crotchet prevented, except in thoſe Baſons that are ſo narrow, that it is impoſſible to deliver without diminiſhing the Bulk of the Head.

Vide Table XXXIX. alſo Vol. I. Lib. III. Chap. 4. Sect. 5. Vol. III. Coll. 34, 35.

THE

THIRTY-SIXTH TABLE

Repreſents, in a lateral View of the *Pelvis*, the Method of extracting with the Aſſiſtance of a curved Crotchet, the Head of the *Fœtus*, when left in the *Uterus*, after the Body is delivered and ſeparated from it; either by its being too large, or the *Pelvis* too narrow.

A.B.C The *Os Sacrum* and *Coccyx*.
D The *Os Pubis* of the left Side.
E.E The *Uterus*.
F The locking part of the Crotchet.
g.h.i The point of the Crotchet on the Inſide of the *Cranium*.

If this Caſe happens from the Forehead's being towards the *Pubes*, or the Child long dead, and ſo mortified that both the Body and Under-jaw are ſeparated unexpectedly, the long Forceps that are curved upwards will be ſufficient to extract the Head; but if the ſame is large, and the *Pelvis* narrow, and the Delivery cannot be effected by the above Method, then the Head muſt be opened, that its Bulk may diminiſh, as it is extracted. The Patient being placed either on her Back, or Side, as in the Explanation of Table XVI. and XXIV. the left hand of the Operator is to be introduced into the *Uterus*, and the Forehead of the *Fœtus* turned to the right Side of the Brim of the *Pelvis*, and a little backwards, the Chin being downwards; after which the palm of the hand and fingers are to be advanced as high as the *Fontanelle*, and the Head graſped with the thumb and little finger on each ſide, as firm as is poſſible, whilſt an aſſiſtant preſſes on each ſide of the *Abdomen* with both hands, to keep the *Uterus* firm in the middle and lower part of the ſame. This done, the Operator having with his right hand introduced and applied the Crotchet to the Head, (the point being turned towards the Forehead, and the convex part towards the *Sacrum*) he muſt go up along the inſide of the left hand as high as the *Fontanelle*, and there, or near it, fix the point of the Crotchet, keeping ſtill the left hand in the former poſition, 'till with the other he pierces the *Cranium* with the point of the Inſtrument, and tears a large opening in it from K to I after this, keeping the Crotchet ſteady, he may ſlide down his left hand in a cautious manner, leſt the former poſition ſhould be altered, and the Head will ſink lower down by the aſſiſtant's preſſing on the *Abdomen*. The two fore-fingers of the left hand are then to be introduced into the Mouth, and the thumb below the Under-jaw, the hand being above the blade of the Crotchet. When this firm hold is taken, the Operator may begin and pull ſlowly with both hands, and as the Brain diſcharges thro' the perforation, the Head will diminſh, and come along. If this Method ſhould fail from the Slipperineſs of the Head, or it's being ſo much oſſified that a ſufficient opening cannot be made, the *Vertex* muſt be turned down to the Brim of the *Pelvis*, the *Fontanelle* backwards, and each blade of the long Forceps introduced along the ſides of the Head, with the curved ſide towards the *Pubes*. After they are joined and locked, the handles are to be tied together with a fillet, to keep them firm on the Head, an aſſiſtant is to keep the handles backwards till the *Cranium* is largely opened with the long Sciſſars ſhewn in Table XXXIX. This done, the Head is to be extracted in a ſlow manner, firſt turning the Forehead to the ſide of the Brim, and as the Brain evacuates, and the Head comes lower down, again turning the Forehead into the Concavity of the *Sacrum*, and completing the Delivery, as in Table XVI.

This Table may alſo ſerve for an Example, to ſhew the Method of fixing the Crotchet on the Head, when altho' the Body is not ſeparated from it, yet it cannot be delivered with the Operator's hands, or the long Forceps, as in Table XXIX. and XXXV.

Vide Vol. I. Lib. III. Chap. 3. Sect. 7. Chap. 4. Sect. 5. alſo Vol. III. Coll. 31, 36.

A
B
C
D
E
E
F
g
h
i
k


a
b
c

THE

THIRTY-SEVENTH TABLE,

With the two following, repreſents ſeveral kinds of Inſtruments uſeful in laborious and difficult Caſes.

A The ſtreight ſhort Forceps, in the exact proportion as to the width between the Blades, and length from the points to the locking part: the firſt being two and the ſecond ſix Inches, which with five Inches and a half, (the length of the handles) makes in all eleven Inches and a half. The length of the handles may be altered at pleaſure. I find however in practice that this ſtandard is the moſt convenient, and with leſs difficulty introduced, than when longer, having alſo ſufficient force to deliver in moſt Caſes, where their aſſiſtance is neceſſary. The handles and loweſt part of the Blades may as here be covered with any durable Leather, but the Blades ought to be wrapped round with ſomething of a thinner kind, which may be eaſily renewed when there is the leaſt ſuſpicion of venereal Infection in a former Caſe: by being thus covered, the Forceps have a better hold, and mark leſs the Head of the Child. For their eaſier Introduction, the Blades ought likewiſe to be greaſed with Hogs-lard.

B Repreſents the poſterior part of a ſingle Blade, in order to ſhew the width and length of the open part of the ſame, and the form and dimenſions of the whole. The handles however as here repreſented are rather too large.

Vide Table XXI. for the figure and dimenſions of the long Forceps, that are curved upwards, and covered in the ſame manner as the former.

The Forceps were at firſt contrived to ſave the *Fœtus*, and prevent, as much as poſſible, the uſe of ſharp Inſtruments; but even to this ſalutary method recourſe ought not to be had but in Caſes where the degree of force requiſite to extract will not endanger by it's conſequences the life of the mother. For, by the imprudent uſe of the Forceps, much more harm may be done than good.

See the Explanation of Table XVI. alſo the Preface to Vol. II. with the Caſes in the Collection on that Subject.

C The Blunt Hook which is uſed for three purpoſes.

Firſt, To aſſiſt the extraction of the Head after the *Cranium* is opened with the Sciſſars, by introducing the ſmall end along the Ear on the outſide of the Head to above the Under-jaw, where the point is to be fixed; the other extremity of the Hook being held with one hand, whilſt two fingers of the other are to be introduced into the foreſaid opening, by which holds the Head is to be gradually extracted.

Secondly, The ſmall end is uſeful in abortions in any of the firſt four or five Months to hook down the *Secundines*, when lying looſe in the *Uterus*, when the Patient is much weakened by Floodings from the too long retention of the ſame, the pains alſo being unable to expel them, and when they cannot be extracted with the fingers. But if the *Placenta* ſtill adheres it is dangerous to uſe this or any other Inſtrument to extract the ſame, as it ought to be left 'till it ſeparates naturally. If a ſmall part of the *Secundines* is protruded thro' the *Os Uteri*, and pulled away from what ſtill adheres in the *Uterus*, the mouth of the *Womb* contracts, and that irritation is thereby removed which would have continued the pains, and have ſeparated and diſcharged the whole.

Thirdly, The large Hook at the other end is uſeful to aſſiſt the extraction of the Body, when the Breech preſents, but ſhould be uſed with great caution, to avoid the diſlocation or fracture of the Thigh.

Vide Table XXIX. alſo Vol. I. Lib. II. Chap. 3. Lib. III. Chap. 3. Sect. 7. and Chap. 4. Sect. 2. Vol. II. Coll. 12. Vol. III. Coll. 31, 32.

THE

THIRTY-EIGHTH TABLE.

A Repreſents the Whale-bone Fillet which may be ſometimes uſeful in laborious Caſes, when the Operator is not provided with the Forceps in ſudden and unexpected Exigencies.

When the *Vertex* of the *Fœtus* preſents, and the Head is forced down into the lower part of the *Pelvis*, the Woman weak and the pains not ſufficient to deliver it, the double of the Fillet is to be introduced along the fore-part of the *Parietal* Bones to the Face, and if poſſible above the Under-jaw; which done, the Whale-bone may be either left in or pulled down out of the Sheath, and every weak Pain aſſiſted by pulling gently at the Fillet. If the Head can be raiſed to the upper part of the *Pelvis*, the Fillet will be more eaſily got over the Chin, which is a ſafer and better hold, than on the Face. If the Face or Forehead preſents, the Fillet is to be introduced over the *Occiput*. *Vide* Vol. I. Lib. III. Chap. 3. Sect. 2. Vol. II. Coll. 24.

In ſuch Caſes likewiſe the Whale-bone may be ſupplied by a Twig of any tough Wood, mounted with a limber Garter or Fillet ſewed in form of a long Sheath.

B.B Gives two Views of a new kind of Peſſary for the *Prolapſus Uteri*, being taken from the *French* and *Dutch* kind. After the *Uterus* is reduced, the large End of the Peſſary is to be introduced into the *Vagina*, and the *Os Uteri* retained in the concave part, where there are three holes to prevent the ſtagnation of any moiſture. The ſmall End without the *Os Externum* has two tapes drawn thro' the two holes, which are tied to four other tapes, that hang down from a belt that ſurrounds the Woman's Body, and by this means keep up the Peſſary. This ſort may be taken out by the Patient when ſhe goes to Bed, and introduced again in the morning; but as this ſometimes rubs the *Os Externum*, ſo as to make it's uſe uneaſy, the round kind marked C. are of more general uſe. They are made of Wood, Ivory, or Cork, (the laſt covered with Cloth and dipt in Wax:) the Peſſary is to be lubricated with Pomatum, the edge forced thro' the paſſage into the *Vagina*, and a finger introduced in the hole in the middle lays it acroſs within the *Os Externum*. They ought to be larger or ſmaller, according to the wideneſs or narrowneſs of the paſſage, to prevent their being forced out by any extraordinary ſtraining. *Vide* Vol. I. Lib. IV. Chap. 1. Sect. 7. Vol. III. Coll. 24.

D.D Gives two Views of a Female Catheter to ſhew it's degree of curvature and different parts. Thoſe for common uſe may be made much ſhorter for conveniency of carrying in the pocket; but ſometimes, when the Head or Body of the Child preſſes on the Bladder above the *Pubes*, it requires one of this length; and in ſome extraordinary Caſes I have been obliged to uſe a Male Catheter.

Vide Vol. I. Lib. II. Chap. 1, 2. Vol. II. Coll. 10. No. 2.

B

A



B

C

D



D




c
a
b
d

THE

THIRTY-NINTH TABLE.

a Repreſents a pair of curved Crotchets locked together in the ſame manner as the Forceps: It is very rare that the uſe of both is neceſſary, excepting when the Face preſents with the Chin turned to the *Sacrum*, and when it is impoſſible to move the Head to bring the Child footling, or deliver with the Forceps. In that caſe, if one Crotchet is not ſufficient, the other is to be introduced, and when joined together will act both as Crotchets, in opening the *Cranium*, and as the Head advances, will likewiſe act as Forceps in moving and turning the Head more conveniently for the Delivery of the ſame. They may alſo be uſeful to aſſiſt when the Head is left in the *Uterus*, and one Blade is not ſufficient. There is ſeldom occaſion however for the ſharp Crotchet, when the Head preſents, the blunt Hook in Table XXVII. being commonly ſufficient, or even the Forceps to extract the ſame, after it is opened with the Sciſſars. Great care ought to be taken when the ſharp Crotchet is introduced, to keep the point towards the *Fœtus*, eſpecially in Caſes where the Fingers cannot be got up to guide the ſame. The dotted lines along the inſide of one of the Blades, repreſent a Sheath that is contrived to guard the point 'till it is introduced high enough; the ligature at the handles marked with the two dotted lines is then to be untied, the ſheath withdrawn, and the point, being uncovered, is fixed as directed in Table XXXVI.

The point, guarded with this Sheath, may alſo be uſed inſtead of the blunt Hook.

b Gives a View of the back-part of one of the Crotchets which is twelve Inches long.

c Gives a front View of the point to ſhew it's length and breadth, which ought to be rather longer and narrower than here repreſented.

d Repreſents the Sciſſars proper for perforating the *Cranium* in very narrow and diſtorted *Pelvis's*. They ought to be made very ſtrong, and nine Inches at leaſt in length, with ſtops or reſts in the middle of the Blades, by which a large dilatation is more eaſily made.

The above Inſtruments ought only to be uſed in the moſt extraordinary Caſes, where it is not poſſible to ſave the Woman without their aſſiſtance.

Vide Vol. I. Lib. III. Chap. 3. Sect. 5. Chap. 5. No. 1. Vol. III. Coll. 31. 35.

ERRATA.

Table IV. Line 19. read Secondly
—— VI. l. 24. r. Fœtus
—— XII. l. 21. r. Labium
—— XXII. l. 1. r. Fœtus
—— XXVI. l. 1. r. out